REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)

2. REPORT DATE 05/06/77 3. REPORT TYPE AND DATES COVERED

4. TITLE AND SUBTITLE
DETERMINATION OF DECONTAMINATION CRITERIA, DIMP AND DCPD (U)

5. FUNDING NUMBERS


6. AUTHOR(S)

O'DONOVAN, P.

DAMD 17 C 5069

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

AEROJET ORDNANCE & MANUFACTURING COMPANY DOWNEY, CA



8. PERFORMING ORGANIZATION REPORT NUMBER

81320R21

9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)

10. SPONSORING / MONITORING AGENCY REPORT NUMBER

FORT DETRICK (FREDERICK, MD.)
FREDERICK, MD

11. SUPPLEMENTARY NOTES

19950119 015

12a. DISTRIBUTION / AVAILABILITY STATEMENT

12b. DISTRIBUTION CODE

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION IS UNLIMITED

13. ABSTRACT (Maximum 200 words)

THE PROPOSAL IS SUBMITTED IN RESPONSE TO A VERBAL REQUEST FROM THE U.S. ARMY MEDICAL RESEARCH & DEVELOPMENT COMMAND. THE PROPOSED PROGRAM IS A CONTINUATION OF THE WORK BEGUN UNDER CONTRACT DAMD-17-75-C-5069. A PORTION OF THE PROPOSED EFFORT IS UNINTERRUPTED CONTINUATION STUDIES OF EXPERIMENTS CURRENTLY UNDERWAY (E.G., PLANT GROWTH & DIMP LYSIMETER TYPE) & THE REMAINING PORTION IS WORK TO BE INITIATED (E.G., DCPD LYSIMETER STUDIES). CURRENTLY A SELECT GROUP OF PLANTS ARE BEING EXPOSED TO DIMP & DCPD IN A SOIL CULTURE SYSTEM TO DETERMINE THEIR EFFECTIVE LEVELS AS TO PHYTOTOXICITY & BIOACCUMULATION IN THIS GROWTH MEDIUM. COMPLETION OF THIS TASK, AS WELL AS THE UTILIZATION OF SEEDS FROM THIS TASK IN FURTHER STUDIES, ARE INCLUDED IN THE PROPOSED EFFORT. A SECOND AREA OF STUDY CURRENTLY UNDERWAY, & TO BE CONTINUED, RELATES TO THE MANNER IN WHICH THE CONTAMINANT COMPOUNDS MIGRATE THROUGH SOIL.

Tere courted executions

14. SUBJECT TERMS

CONTAMINANTS, FLORA, SOIL, CHEMICALS, LYSIMETER

15. NUMBER OF PAGES

16. PRICE CODE

17. SECURITY CLASSIFICATION OF REPORT

OF THIS PAGE

SECURITY CLASSIFICATION

19. SECURITY CLASSIFICATION
OF ABSTRACT

20. LIMITATION OF ABSTRACT

AEROJET ORDNANCE AND MANUFACTURING COMPANY 9236 East Hall Road Downey, California 90241

DETERMINATION OF DECONTAMINATION CRITERLA

DIMP AND DCPD (U)

Rocky Mountain Arsenal Information Center Commerce City, Colorado

Report No. 1953-01(21)MP

Contract DAMD-17-C-5069

| Accesio | n For | |
|-------------------------------------|----------------|-------------------|
| NTIS DTIC Unanno Justifica | TAB ounced | X |
| By Distribu | ıtion / | |
| A | vailabilit | y Codes |
| Dist | Avail a Spe | and / or ecial |
| A-1 | | |

To

U. S. Army Fort Detrick Fredrick, Maryland 21701

| FILE CO | \mathbf{OPY} | |
|---------|----------------|--|
|---------|----------------|--|

| Prepared by: P. A. O | O'Nonovan Donovan |
|----------------------|----------------------|
|----------------------|----------------------|

| Date: | 6 | May | 1977 | |
|-------|------|-----|------|--|
| | | | | |
| | | | | |
| | | | | |

| ** - | | D | 19 |
|------|----|--------|-----|
| NO. | OI | Pages: | T-7 |
| 110. | - | ~ =- 5 | |

| | DESCRIPTION | E | - S | 0 | 2 | ٥٠ | u. | 3 | ₹ | 2 | : | <u> </u> | 5 | 0 | z: | 0: | <u> </u> | - | \\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\ | 3 |
|----------|--|---|----------|------------------|-----|----|----|-----|---|----------|----------|------------|----------|----|----|----|----------|---|--|-----|
| " | SURVEY OF LITERATURE | | 7 | 1 | s . | 8 | - | 6 | | | 2 | | 2 | 2 | -# | | | | -11 | × × |
| Ď., | PROTOCOL TASK II HYDROPONIC EXPERIMENTS | | <u> </u> | | | | | | | | | | | | | | | | | |
| | SELECT PLANTS INSTALL APPARATUS | | ┰╂ | — _Y - | | | | | | | | | | | | | | | | |
| | GERMINATE SEEDS GROW AND INOCULATE PLANTS PHOTOGRAPHIC AND CHEMICAL ANALYSIS | | 11. | | | | -H | | | 77 | r | | | | | | | | | |
| 울. | PROTOCOL TASK III (PART 1) | | - | _ | | | - | | | \vdash | ├— | <u> -</u> | _ | _ | | - | - | | | |
| 6 | SOIL CULTURE EXPERIMENTS CONSTRUCT GREENHOUSE | | | • | | | | - 1 | | | | | | | | | | | | |
| | PREPARE TEST PLAN | | | | | | | | 1 | | | | | | | | | | | |
| | GROW AND INOCULATE PLANTS | | | | | | | | | 1 | 十 | ╂ | ŀ | į. | | | <u></u> | - | | |
| | PHOTOGRAPHIC AND CHEMICAL ANALYSIS | - | | | | | | | | | 1 | -{ | _ | Ų | | | \vdash | - | _, | |
| | RADIOACTIVE DCPD TRACING | | | | | | | | | | | | | | | 1 | + | - | | |
| - | LYSIMETER STUDIES | | \vdash | \vdash | ļ | | - | _ | | | | | <u> </u> | | | | | | | |
| | PROCURE, PROCESS AND FABRICATE LYSIMETERS | | | - | | I | | | | | | | | | | | | | | |
| | CHRONIC DIMP | | | | | | - | - | 1 | | 1 | ┪ | -} | - | | | | | _ | |
| | SINGLE CHARGE DIMP | | | | | | | | 1 | 1 | 1 | + | - | Ļ | I | 1 | + | + | 1 | |
| | DEVELOP ANALYSIS FOR DCPD IN SOIL | | H | H | Ц | | H | | | | \vdash | | | | | | \dashv | - | | |
| ۵ | DATA | | | · | | | | | | | | | | | | | | | | |
| • | ANNUAL REPORT | | | | | | | | | | | | _1 | -1 | | | | | | |

▼ - Satisfactory Progress - on schedule

Determination of Decontamination Criteria - DIMP and DCPD Research Task Schedule

Progress on items proposed for action during April, 1977 is discussed in the following paragraphs.

Full Scale Lysimeter Tests

All of the lysimeter irrigation experiments have been terminated and the final multiple core samples are being analyzed for DIMP content. The lysimeters were divided into two groups, which consisted of five lysimeters each. The five lysimeters were filled, one each, with the following types of soil:

1. Chino - sandy clay loam

2. Brawley - silty clay

3. Ventura - clay loam

4. Fullerton - sandy loam

5. Walnut - clay loam

Group 1 was subjected to irrigation on a regular basis with 20 ppm (parts per million) DIMP (disopropyl methyl phosphonate). Group 2 had 20 ppm DIMP intimately mixed with the top 1 foot layer of soil and subsequently irrigated with regular additions of distilled water. The general trend in DIMP concentrations in group 1 core samples has been for the surface layer to be relatively high and for the successively deeper layers to rapidly fall off in concentration. The multiple samples analyzed here and in group 2 will give a more significant basis for the reliability of these concentration profiles.

The raw data from the group I terminal samples are shown in Table 1.

Taking the average of each of the sets of figures from Table 1 gives the more concise data in Table 2. These data are plotted in Figures 1 through 5 and illustrate the above stated trends.

The group 2 lysimeter soils have been sampled and are currently undergoing analysis. In this case as in Group 1 cores were taken from four separate

Table 1
DIMP Content of Soil Jamples (ppm) 426 Days

| | | | | | | | | | | | | | | • |
|---------|-----------|-----|---------------------|----------|---------|----------|----------|----------|----------|----------|----------|----------|----------|-----|
| | | (4) | 34.4 | 5.7 | 11.6 | 5.7 | 8.1 | 5.6 | 8.0 | 5.4 | 8.3 | 5.4 | 7.0 | |
| | EY | (3) | 15.8 | 7.8 | 5.6 | 9.1 | 9.7 | 8. | 7.2 | 7.3 | 6.4 | 5.9 | 7.5 | |
| | BRAWLEY | (2) | 8.6 | 5.9 | * | 6.4 | 8.0 | 8.9 | 4.8 | 5.2 | 3.7 | 4.2 | 4.2 | |
| | В | (1) | 14.8 | * | * | 6.9 | 4.5 | 6.4 | 6.2 | 5.0 | 5.7 | 4.3 | 4.3 | |
| | | (4) | 17.5 | 9.1 | 9.7 | 12.7 | 8.1 | 10.2 | 12.0 | 9.5 | 13.6 | 14.3 | 8.6 | |
| | T | (3) | 40.3 | 4.6 | 11.1 | 5.9 | 7.4 | 5.3 | 7.5 | 6.3 | 11.5 | 7.5 | 4.9 | |
| | WALNUT | (2) | 26.2 | 6.2 | 5.2 | 3.8 | 3.8 | 5.1 | 5.1 | 4.1 | 4.4 | 4.2 | 7.4 | |
| T BL | Marie 1 | (1) | 49.0 | 16.2 | 6.9 | 6.2 | 4.5 | 5.4 | 5.5 | 6.7 | 5.5 | 5.2 | 4.1 | |
| | ¢. | (4) | 34.9 | 10.9 | 12.0 | 7.9 | 10.4 | 9.1 | 10.5 | 7.3 | 7.3 | 8.9 | 5.1 | |
| | ron | (3) | 14.3 | 13.2 | 5.6 | 7.1 | 5.9 | 0.9 | 13.6 | 7.9 | 9.6 | 12.3 | 6.9 | |
| | FULLERTON | (2) | 23.7 | 3.9 | 3.9 | 3.1 | 3.1 | 3.3 | 2.9 | 2.0 | 2.4 | 3.4 | 2.5 | |
| Group 1 | FI | Œ | 21.3 | 8.9 | 6.8 | 6.3 | 4.0 | 4.4 | 6.2 | 0.9 | 5.1 | 3.1 | 5.1 | |
| Gre | | (4) | 40.0 | 7.7 | 8.3 | 5.3 | 7.7 | 4.4 | 4.5 | 4.4 | 3.4 | 4.2 | 4.3 | |
| | | (3) | 10.0 | 5.9 | 9.9 | 4.4 | 4.7 | 5.4 | 5.3 | 5.4 | 4.3 | 5.3 | 5.4 | |
| | CHINO | (2) | 27.4 | 7.4 | 7.1 | 6.1 | 3.8 | 6.4 | 1.2 | 1.7 | 1.6 | 2.0 | 10.6 | |
| | Ĭ | (1) | 38.3 | 8.5 | 6.4 | 5.5 | 4.6 | 3.4 | 3.0 | 6.4 | 2.6 | 2.6 | * | |
| | | (4) | | 5.7 | 8.5 | 4.0 | 5.3 | 3.6 | 5.0 | 11.5 | 4.0 | 6.2 | 5.1 | |
| | ≴ | (3) | 24.7 | 11.3 | 3.9 | 3.0 | 2.7 | 1.9 | 2.8 | 2.5 | 3.1 | 6.0 | 1.5 | |
| - | VENTURA | (2) | 22.4 | 3.1 11.3 | 3.0 | 1.5 | 2.1 | 2.6 | 2.2 | 2,3 | 2.6 | 2.3 | 3.7 | |
| | | (E) | 22.0 22.4 24.7 44.5 | 5.7 | 3.8 | 1.5 | 3.2 | 1.4 | 0.8 | 1.6 | 1.7 | 1.7 | 2.0 | |
| | Depth | | 0 (surface) | 1.9 - 0 | 6 – 12" | 12 – 18" | 18 – 24" | 24 - 30" | 30 - 36" | 36 – 42" | 42 - 48" | 48 - 54" | 54 - 60" | |
| | 1 | 1 | | | | | | | | _ | | | | - 1 |

*<0.1 ppm


Table 2


Average DIMP Content of Soil Samples (ppm)

426 days, Group 1


| Depth | Ventura | Chino | Fullerton | Walnut | Brawley |
|-------------|---------|-------|-----------|--------|---------|
| | | | | | |
| 0 (surface) | 28.4 | 28.9 | 23.6 | 33.3 | 18.4 |
| 0-6" | 6.5 | 7.4 | 8.7 | 9.0 | 6.5* |
| 6-12" | 4.8 | 7.1 | 7.1 | 8.2 | 8.6* |
| 12-18" | 2.5 | 5.3 | 6.1 | 7.2 | 7.0 |
| 18-2411 | 3.3 | 5.2 | 5.9 | 6.0 | 7.6 |
| 24-30" | 2.4 | 4.9 | 5.7 | 6.5 | 6.9 |
| 30-36" | 2.7 | 3.5 | 8.3 | 7.5 | 6.6 |
| 36-42" | 4.5 | 4.1 | 5.8 | 6.6 | 5.7 |
| 42-48'' | 2.9 | 3.0 | 6.1 | 8.8 | 6.0 |
| 48-54" | 2.8 | 3.5 | 6.4 | 7.8 | 5.0 |
| 54-60" | 3.1 | 6.8* | 4.9 | 6.3 | 5.8 |
| | | | | | |

^{*} Group contains some samples with no detectable DIMP i.e., < 0.1 ppm.






Sample Depth (in)
Figure 2. DIMP Content of soil samples, Brawley, at 426 days,
Group I




Sample Depth (in)
Figure 3. DIMP Content of soil samples, Ventura, at 426 days,
Group I.



DIMP Conte

Figure 4. DIMP Content of soil samples, Fullerton, at 426 days, Group I.



DIMP Conter (ppm)

Figure 5. DIMP Content of soil samples, Walnut, at 426 days, Group I.

locations in each lysimeter. This data will be included in a future report.

Soil Culture Experiments

Harvesting of the broad range (0-1000 ppm) soil growth tests is completed. Data on the biomass of the sugar beet, alfalfa, carrot and bean are given in tables 3, 4, 5 and 6. Plotting the mass data for the normally edible portion of the plants gives the graphs shown in figures 6, 7, 8 and 9. In general this data shows that for sugar beets and alfalfa 50 ppm DIMP results in poorer than control yields and for carrots and beans somewhere between 50 and 100 ppm DIMP gives poorer than control yields. In all cases the yields of DCPD treated plants are significantly higher than the DIMP plants and generally higher than the control plants.

Radioactive Tracing of Soil Contamination

Further data is available from the radioactive tracer experiments described in 1953-01(18)MP. The most recent data is shown in Table 7. This data confirms the preliminary data in 1953-01(19)MP which generally indicated that passing air over columns of day soil impregnated with either radioactive DIMP or DCPD, at 20 ppm, does not significantly lower the concentration of radioactivity in the soil.

Similar experiments using moistened soil conditions and larger diameter soil containers are currently underway and data from these tests should be available in a future report.

Table 3

Tabulation of Average Weight of Plant Parts @ 201 Days of Age

| | Ave | erage V | V e i g h t | (g) of | | Number of | Contaminant | |
|---------------|------------------|----------|-------------|--------------------------|------------------|-------------------------|-------------|-----------------------|
| Plant Type | Leaf | Stem | Root | Edible Root/ Plant | Total Plant | Plants in Average | Type | H ₂₀ (ppm) |
| Sugar | 1// 02 | | | 187.35 | 354.28 | 4 | DIMP | Control |
| Beet " | 166.93 | - - | - | 12.00 29.50 | 28.68 63.58 | 5 | | 50 100 |
| 11 19 2 | 34.08 20.74 | - | - , | 6.38 | 27. 12 | _ 5 | 11.7 | 300 |
| ** | 1.05 94.30 | - | - | 1.51 | 140.90 | 5 4 | DCPD " | Control |
| 11 | 115.13 | - | - | 112.38 | 227.51 309.44 | 3 | 11 | 100 |
| ar ir | 161.93 102.43 | - | - | 188.37 78.25 | 350.30 180.68 | 4 | 11 | 500 |
| 11 | 85.45 133.05 | - | - | 91.50 | 176.95 268.75 | 2 2 | 11 | 1000 |
| • | | | | | | | | |
| | | | | | <u> </u> | l | | |

Table 4

Tabulation of Average Weight of Plant Parts @ 201 Days of Age

| | Αve | erage V | Veight | (g) of | | Number of | Contaminant | 1 |
|---------------|------|---------|--------|--------------------------|----------------|-------------------------|-------------|-----------------------|
| Plant Type | Leaf | Stem | Root | Edible Root/ Plant | Total Plant | Plants in Average | Type | H ₂₀ (ppm) |
| Alfalfa | 4.89 | 8.68 | 4.25 | S 🛶 . S | 17.82 | 15 | DIMP | Control |
| 11 | 2.72 | 5.19 | 2.24 | - | 10.15 | 14 | tt | 50 |
| 11 | 0.92 | 4.20 | 1.19 | - | 6.31 | - 11 | it . | 100 |
| iris s | 0.60 | 1.03 | 0.12 | | 1.75 | 3 | 11 | 300 |
| 11 | 1.48 | 3.39 | 0.22 | - | 5.09 | 2 | 11 | 500 |
| 11 | 4.20 | 4.74 | 3.24 | _ | 12.18 | 12 | DCPD | Control |
| it · | 6.35 | 10.58 | 4,45 | - | 21.38 | 12 | in) | 50 |
| 11 | 8.84 | 12.44 | 6.04 | .= | 27.32 | 7 | . 11 | 100 |
| 11 | 4.94 | 9.72 | 3.92 | - | 18.58 | 5 | · tt | 300 |
| 11 | 6.36 | 14.09 | 4.37 | _ | 24.82 | 2 | 11 | 500 |
| | a. | | | | | | | |
| | | | · | | | | | |
| · | | | | • | | | | |
| | : . | | 1 | | | | | |
| | | | | | | | | |
| | I | | 1 | I | 1 | 1 | ı | 1 |

The 700 and 1000 ppm plants did not survive the experiment.

Table 5

Tabulation of Average Weight of Plant Parts @ 201 Days of Age

| | · | | | | | | | Conc. of |
|--|----------------------|----------------------|----------|--------------------------|----------------|-------------------------|-------------|-----------------------|
| | Αve | erage V | Veight | (g) of | | Number of | Contaminant | Contami- nant in |
| Plant Type | Leaf | Stem | Root | Edible Root/ Plant | Total Plant | Plants in Average | Type | H ₂₀ (ppm) |
| Carrot | 1.29 | 1.54 | - | 17.01 19.65 | 19.84 21.73 | 9 | DIMP | Control |
| 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 | | 1.33 0.50 0.84 | | 0.05 0.40 | 0.72 A | 2 | tt tt | 300 500 |
| 11 | 2.19 2.49 7.80 | 3.18 11.40 | - | 17.49 39.40 | 23.16 | 8 | DCPD " | Control |
| 11 | 5.37 3.87 | 10.03 9.58 | - | 50.57 63.23 | 65.97 76.68 | 3 6 | 11 | 300 500 |
| n . | 4.30 4.42 | 7.35 6.40 | - | 36.65 50.70 | 48.30 61.52 | 2 4 | 11 | 1000 |
| | | | | | | | | |
| | | | | | | | | |

Table 6

Tabulation of Average Weight of Plant Parts @ 201 Days of Age

| Plant Type | A v d | erage V Stem | Veight Root | (g) of Edible Fruit/ Plant | Total Plant | Number of Plants in Average | Contaminant Type | Conc. of Contami- nant in H ₂₀ (ppm) |
|---------------|-------|-----------------|----------------|-------------------------------------|----------------|---|---------------------|--|
| | | | | I lailt | | | | |
| Bean | 2.13 | 9.83 | 1.17 | 9.67 | 22.80 | . 3 | DIMP | Control |
| 11 | 3.00 | 34.50 | 7.30 | 22.30 | 67.10 | 1 | 11 | 50 |
| | 9.10 | | 1.85 | 4.95 | 30.90 | 2 | 11 | 100 |
| | 25.30 | 29.80 | 14.90 | 21,50 | 91.50 | 1 | DCPD | 100 |
| 11 | 51.20 | 35.70 | 30.5 | 17.40 | 134.80 | 1 | 11 | 300 |
| 11 | 4.50 | 4.85 | 2.40 | 14.00 | 25.75 | 2 | . 11 | 500 |
| . 11 | 7.95 | 17.55 | 13.60 | 13.25 | 52.35 | 2 | 11 | 700 |
| ft | 4.85 | 13.00 | 8.05 | 13.65 | 39.55 | 2 | 11 | 1000 |
| \ \ | | | | | | | | |
| | | | | | | · | | |
| | | | | | | | | |




Figure 6. Average Yield of Sugar Beets Irrigated with DIMP or DCPD Contaminated Water. 14




Figure 7. Average Yield of Alfalfa Irrigated with DIMP or DCPD Contaminated Water. 15





Figure 8. Average Yield of Carrots Irrigated with DIMP or DCPD Contaminated Water.



Concentration of Contaminant (ppm)

Figure 9. Average Yield of Beans Irrigated with DIMP or DCPD

Contaminated Water.

Table 7

| Scintillation Count Data From DIMP/DCPD Tracer Experiments in Fullerton Soil | | | | | | |
|--|-------------------|---------------------|---|--|-----------------------|--|
| Sample No. | Ex Description | Sample wt (g) | s in Fullerto Radioact Calculated # Ci | lvity | % of Stock Rad. Re | Air flow time @ 100 ml/min (hrs) |
| 1 | Stock DCPD Soil | 5.0098 | 8x10 ⁻³ * | 7.86×10 ⁻³ | 100 | |
| 2 | Stock DIMP Soil | 6.1420 | 3.37x10 ⁻¹ * | 3.12×10 ⁻⁴ | 100 | |
| 3 | 0-1" Soil DCPD | 11.9831 | | 0.18 | | |
| 4 | 1"-2" " | 13.1534 | ************************************** | 0.18 | | 267 |
| 5 | 2"-3" " | 12/3646 | in 1 | $\begin{vmatrix} 0.14 \\ 0.65 \\ 0.14 \end{vmatrix}$ | | 201 |
| 6 | 3"-4" " DCPD Trap | 16,2240 12,4104 | 0.4 | 0.01 | 165 | |
| 11 7 | 0-1" Soil DIMP | 13. 9275 | 0, 1 | 4.14 | | : |
| 8 | 1"-2" " | 15.3755 | · | 5.00 | | |
| 9 | 2-3" | 10.3126 | | 3.20 | | 231 |
| 10 | 3-411 11 | 11.0975 | | 3.82 | |) |
| 123 | DIMP Trap | 17.5540 | 16.9 | 0.03 | 104 | |
| 13, | DIMP in Soil | 17.1267 | 3.37×10^{-2} | 3.49×10 ⁻ * | 112 | |
| 14 | DCPD in Soil | 17.2577 | 8x10 ⁻³ * | 7.90×10 | 101 | |
| 16 | Dil. Stock DCPD | 0.1027 | 3.3x10 ² * | | 102 | |
| | | | | | | |
| | | | | | | |
| | | | | | | |

#Ci/g

PROPOSED ACTIVITY FOR MAY 1977

- o Terminate all experimental activities including final anlayses of samples of both soil and plants.
- o Begin preparation of draft copy of project final report.